

TYLOSIN 20% Injectable solution

COMPOSITION (per ml)

Tylosin base (as tartrate)..... 200mg

TARGET SPECIES

Cattle, lambs, pigs, dogs and cats

INDICATIONS

For the treatment of infections caused by bacterias sensitive at tylosine and especially for dealing infections caused by mycoplasmas.

DOSAGE AND METHOD OF ADMINISTRATION

The recommended dosage is 4 – 10 mg per kg of body weight

- Cattle: 0,2-0,5 ml per 10 kg of body weight, daily, for 3-5 days. Do not inject at the same spot quantity of solution greater than 10 ml. If a big animal requires p.e 30 ml , then 3 injections of 10 ml at 3 different spots should be done.
- Lambs: 0,5 ml / 10 kg of body weight for 3-5 days
- Pigs: 0,1-0,5 ml per 10 kg of body weight daily , for 3-5 days. Do not inject solution more than 5 ml at the same spot
- Dogs and cats: 0, 2-0,5 ml per 10 kg of body weight daily, for 3-5 days. Do not inject solution more than 5 ml at the same spot

If within 3 days improvement is not observed, the treatment should be reviewed

CONTRA INDICATIONS

This product should not be administered at horses and rabbits. Also, should not be administer at animals with renal disorders, or hypersensitivity at tylosine

UNDESIRABLE EFFECTS

In some cases, may be observed irritation , light itch, or necrosis at the point of injection. Occasionally, have been referred allergic reactions due to hypersensitivity at tylosine. Rarely, can be observed:

Pigs: edema and proptosis of orthus

Calves and lambs: raise of cardiac frequency and of the rhythm of respiratory movements

Dogs: colpic arrhythmia

INTERACTION WITH OTHER MEDICAMENTS

Do not administer simultaneously with cardiac glycosides, due to the raise of their concentration in blood.

INCOMPATIBILITY

Not mentioned

OVERDOSE

In case of overdose can be caused digestive disorders as vomit, anorexia, diarrhea and also toxicosis. In that case the administration must be stopped at once and the symptoms must be handled.

USE DURING PREGNANCY AND LACTATION

Is not mentioned

SPECIAL PRECAUTIONS FOR USE

Special attention should be given at hyper-development of sensitive micro-organisms and especially of fungus

SPECIAL PRECAUTIONS TO BE TAKEN BY THE PERSON ADMINISTERING THE PRODUCT TO ANIMALS

Avoid direct contact with the product .It is possible to notice dermatitis by contact.

WITHDRAWAL TIME

Meat:	Cattle - pigs :	21 days
	Sheep:	42 days
Sheep milk:		108 hours
Cows milk:		5 days

STORAGE

After the first opening remains stable for 30 days at dry and dark place between 15-25° C

SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIAL, IF ANY:

Not requested

PACKAGE

Flacons of 100 ml.